JOURNAL OF THE AMERICAN PODIATRIC MEDICAL ASSOCIATION

January–December 2001
Volume 91

SUBJECT INDEX

A

Activity
Activity monitors: should we begin dosing activity as we dose a drug? [Armstrong and Boulton], 3:152
Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451

American Indians
The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280

Amputation
Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451
The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445
Differentiation and treatment of phantom sensation, phantom pain, and residual-limb pain [Finnoff], 1:23
Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313
Primary-care interventions for the population at risk for amputation [Yacyshyn and Andrews], 1:2
Rehabilitation after amputation [Esquenazi and DiGiacomo], 1:13
The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533
Subtotal pedal amputations: biomechanical and intraoperative considerations [Funk and Young], 1:6
Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34

Anatomy
The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

Aneurysm
Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93

Ankle
The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Manipulation method for the treatment of ankle equinus (letter) [Menz], 2:105
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356

Antisepsis
Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288

APMA Announcements and Reports
Council on Podiatric Medical Education, 81st annual report, 2000, 3:154

Artery
The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

Arthrodesis
Arthrodesis of the interphalangeal joint of the hallux: a simple and effective technique [Yu et al], 8:427
Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331
Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63

Arthroereisis
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394

Arthropathy
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Hemophilic arthropathy: considerations in management [Chang et al], 8:406
Surgically induced Charcot's foot [Fishco], 8:388

Athletics
Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356
Survey of the effects of aerobic dance on the lower extremity in aerobic instructors [du Toit and Smith], 10:528

Axis of rotation
Scientific approach to the axis of rotation at the midtarsal joint [Nester et al], 2:68

B

β-Glucuronidase
Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114

Balance
Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
Foot pain impairs balance and functional ability in community-dwelling older people [Menz and Lord], 5:222

Benzalkonium chloride
Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288

Biofeedback
Effect of biofeedback-assisted relaxation training on foot ulcer healing [Rice et al], 3:132

JOURNAL OF THE AMERICAN PODIATRIC MEDICAL ASSOCIATION

January–December 2001
Volume 91

SUBJECT INDEX

A

Activity
Activity monitors: should we begin dosing activity as we dose a drug? [Armstrong and Boulton], 3:152
Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451

American Indians
The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280

Amputation
Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451
The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445
Differentiation and treatment of phantom sensation, phantom pain, and residual-limb pain [Finnoff], 1:23
Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313
Primary-care interventions for the population at risk for amputation [Yacyshyn and Andrews], 1:2
Rehabilitation after amputation [Esquenazi and DiGiacomo], 1:13
The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533
Subtotal pedal amputations: biomechanical and intraoperative considerations [Funk and Young], 1:6
Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34

Anatomy
The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

Aneurysm
Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93

Ankle
The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Manipulation method for the treatment of ankle equinus (letter) [Menz], 2:105
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356

Antisepsis
Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288

APMA Announcements and Reports
Council on Podiatric Medical Education, 81st annual report, 2000, 3:154

Artery
The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

Arthrodesis
Arthrodesis of the interphalangeal joint of the hallux: a simple and effective technique [Yu et al], 8:427
Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331
Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63

Arthroereisis
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394

Arthropathy
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Hemophilic arthropathy: considerations in management [Chang et al], 8:406
Surgically induced Charcot's foot [Fishco], 8:388

Athletics
Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356
Survey of the effects of aerobic dance on the lower extremity in aerobic instructors [du Toit and Smith], 10:528

Axis of rotation
Scientific approach to the axis of rotation at the midtarsal joint [Nester et al], 2:68

B

β-Glucuronidase
Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114

Balance
Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
Foot pain impairs balance and functional ability in community-dwelling older people [Menz and Lord], 5:222

Benzalkonium chloride
Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288

Biofeedback
Effect of biofeedback-assisted relaxation training on foot ulcer healing [Rice et al], 3:132

Biomechanics
Changes in foot function with the use of Root and Blake orthoses [Reed and Bennett], 4:184
The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280
The effect of high-Dye and low-Dye taping on rearfoot motion [Keenan and Tanner], 5:255
Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394
Function of the windlass mechanism in excessively pronated feet [Aquino and Payne], 5:245
Manipulation method for the treatment of ankle equinus (letter) [Menz], 2:105
Reduction of peak pressure on the forefoot with a rigid rocker-bottom postoperative shoe [Fuller et al], 10:501
The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121
Reliability of navicular displacement measurement as a clinical indicator of foot posture [Vinicombe et al], 5:262
Scientific approach to the axis of rotation at the midtarsal joint [Nester et al], 2:68
Subtalar joint axis location and rotational equilibrium theory of foot function [Kirby], 9:465
Subtotal pedal amputations: biomechanical and intraoperative considerations [Funk and Young], 1:6
Surgically induced Charcot's foot [Fishco], 8:388
Symmetry of plantar pressures and vertical forces in healthy subjects during walking [VanZant et al], 7:337

Blindness
Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313

Blood
The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Hemophilic arthropathy: considerations in management [Chang et al], 8:406
The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

Bone
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Brachymetatarsia of the third and fourth metatarsals [Goforth and Overbeek], 7:373
The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445
Internal fixation of a displaced tibial sesamoid fracture [Riley and Selner], 10:536
Isolated cuboid fracture: a rare occurrence [Miller and Handzel], 2:85
Isolated gouty tophus of a tibial sesamoid of the foot [Lemont and Sabo], 7:379
Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306
Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369
Posterior facet talocalcaneal coalition [Brekke et al], 8:422
Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142
Stress fracture of the foot secondary to osteoporosis: an atypical presentation [Freeman and Randall], 2:99
Talar fractures: three case studies [Jimenez and Morgan], 8:415

Book reviews
Functional Reconstruction of the Foot and Ankle [Miller], 5:269
Handbook of Lower Extremity Neurology [Miller], 2:102
Textbook of Bunion Surgery [Miller], 5:269

Brachymetatarsia
Brachymetatarsia of the third and fourth metatarsals [Goforth and Overbeek], 7:373

C

Calcaneus
Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369
Posterior facet talocalcaneal coalition [Brekke et al], 8:422
The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121
Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142

Callus
Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275

Charcot's foot
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Surgically induced Charcot's foot [Fishco], 8:388

Children
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394

Cholesterol
Smith-Lemli-Opitz syndrome: a genetic disorder with pedal manifestations [Sheff], 3:149

Coalition, talocalcaneal
Posterior facet talocalcaneal coalition [Brekke et al], 8:422

Cuboid
Isolated cuboid fracture: a rare occurrence [Miller and Handzel], 2:85

D

Dance, aerobic
Survey of the effects of aerobic dance on the lower extremity in aerobic instructors [du Toit and Smith], 10:528

Databases
The case for multiple database searching in podiatric medicine (commentary) [Menz], 2:103

Diabetes mellitus
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451
The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445

The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280
Evaluation of pressure threshold prior to foot ulceration: one-*versus* two-point static touch [Barber et al], 10:508
Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275
Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369
Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313
Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325
A prospective study of predictors for foot ulceration in type 2 diabetes [Kästenbauer et al], 7:343
The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533
Subtotal pedal amputations: biomechanical and intraoperative considerations [Funk and Young], 1:6

Dorsiflexion
Manipulation method for the treatment of ankle equinus (letter) [Menz], 2:105

Dysplasia
Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306

E

Education
A conceptual model for public health education in podiatric medicine [Helfand and Hausman], 9:488
Internet resources for podiatric medical students [Fikar and Fikar], 6:316

Elderly
The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456
Foot pain impairs balance and functional ability in community-dwelling older people [Menz and Lord], 5:222

Enzyme
Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114

Equinus
Manipulation method for the treatment of ankle equinus (letter) [Menz], 2:105
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356

Erythrocyte sedimentation rate
The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445

Ethyl alcohol
Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288

F

Fasciitis, plantar
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356
Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142

Fibrolipoma
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89

Flatfoot
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394

Footprints
A comparison of footprint indexes calculated from ink and electronic footprints [Urry and Wearing], 4:203

Footwear
Reduction of peak pressure on the forefoot with a rigid rocker-bottom postoperative shoe [Fuller et al], 10:501
Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34

Force, vertical
Symmetry of plantar pressures and vertical forces in healthy subjects during walking [VanZant et al], 7:337

Forefoot
Reduction of peak pressure on the forefoot with a rigid rocker-bottom postoperative shoe [Fuller et al], 10:501

Fracture
Internal fixation of a displaced tibial sesamoid fracture [Riley and Selner], 10:536
Isolated cuboid fracture: a rare occurrence [Miller and Handzel], 2:85
Stress fracture of the foot secondary to osteoporosis: an atypical presentation [Freeman and Randall], 2:99
Talar fractures: three case studies [Jimenez and Morgan], 8:415

Fungal infection
The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456
Efficacy of terbinafine for toenail onychomycosis: a multicenter trial of various treatment durations [Pollak and Billstein], 3:127; (letter) [Wallace], 10:542
Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351
Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325
Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540

G

Gait
Changes in foot function with the use of Root and Blake orthoses [Reed and Bennett], 4:184
The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280
Function of the windlass mechanism in excessively pronated feet [Aquino and Payne], 5:245
Symmetry of plantar pressures and vertical forces in healthy subjects during walking [VanZant et al], 7:337

Gangrene
Bilateral forefoot gangrene secondary to Lemierre's disease [Soave and Kuchar], 3:147; (letter) [Kaufman], 7:381

Geriatric population
The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456

Foot pain impairs balance and functional ability in community-dwelling older people [Menz and Lord], 5:222

Gout
Isolated gouty tophus of a tibial sesamoid of the foot [Lemont and Sabo], 7:379

Grading system
The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74

Guidelines, authors'
Guidelines for Authors, 1:50, 5:270, 7:382, 9:496

H

Hallux
Arthrodesis of the interphalangeal joint of the hallux: a simple and effective technique [Yu et al], 8:427
The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74
Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251

Hammer toe deformity
Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63

Hand wash
Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288

Heel
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55
Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356
Posterior facet talocalcaneal coalition [Brekke et al], 8:422
The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121
Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142

Hemangioma
Hemangioma of the foot [Lamm and Pontious], 2:96

Hemophilia
The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Hemophilic arthropathy: considerations in management [Chang et al], 8:406

Human immunodeficiency virus
Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79

Human papillomavirus
Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79

Hyperkeratosis
Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114
Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275

I

Imaging
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
A comparison of radiographic foot measurements taken in two different positions [Bryant], 5:234
Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306

Implant
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394

Indexes
The case for multiple database searching in podiatric medicine (commentary) [Menz], 2:103

Infection
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Bilateral forefoot gangrene secondary to Lemierre's disease [Soave and Kuchar], 3:147
The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456
The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445
Efficacy of terbinafine for toenail onychomycosis: a multicenter trial of various treatment durations [Pollak and Billstein], 3:127; (letter) [Wallace], 10:542
Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351
Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369
Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79
Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325
Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540
Wound healing and infection in nail matrix phenolization wounds: does topical medication make a difference? [Dovison and Keenan], 5:230

Information resources
Internet resources for podiatric medical students [Fikar and Fikar], 6:316

Injury
Internal fixation of a displaced tibial sesamoid fracture [Riley and Selner], 10:536
Isolated cuboid fracture: a rare occurrence [Miller and Handzel], 2:85
Stress fracture of the foot secondary to osteoporosis: an atypical presentation [Freeman and Randall], 2:99
Survey of the effects of aerobic dance on the lower extremity in aerobic instructors [du Toit and Smith], 10:528
Talar fractures: three case studies [Jimenez and Morgan], 8:415

Insoles
The effect of customized insoles on the reduction of post-work discomfort [Sobel et al], 10:515

Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194

Internet
Internet resources for podiatric medical students [Fikar and Fikar], 6:316

J

Joint
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Arthrodesis of the interphalangeal joint of the hallux: a simple and effective technique [Yu et al], 8:427
The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394
Hemophilic arthropathy: considerations in management [Chang et al], 8:406
Internal fixation of a displaced tibial sesamoid fracture [Riley and Selner], 10:536
Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331
Manipulation method for the treatment of ankle equinus (letter) [Menz], 2:105
Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63
Posterior facet talocalcaneal coalition [Brekke et al], 8:422
The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121
Scientific approach to the axis of rotation at the midtarsal joint [Nester et al], 2:68
Subtalar joint axis location and rotational equilibrium theory of foot function [Kirby], 9:465
Surgically induced Charcot's foot [Fishco], 8:388

Journals, podiatric medical
Publication patterns and perceptions of the Australian podiatric medical faculty [Menz], 4:210

K

Keratoderma
Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114

Keratosis
Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114
Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275

L

Laboratory tests
The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445

Laser
Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142

Lateral intermetatarsal angle
Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251

Lemierre's disease
Bilateral forefoot gangrene secondary to Lemierre's disease [Soave and Kuchar], 3:147; (letter) [Kaufman], 7:381

Lesion
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Hemangioma of the foot [Lamm and Pontious], 2:96
Juxtamalleolar lipoma: painful menopausal lipoma [Lemont], 6:311
Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109
Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79
Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93

Lipoma
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Juxtamalleolar lipoma: painful menopausal lipoma [Lemont], 6:311

M

Malleolus
Juxtamalleolar lipoma: painful menopausal lipoma [Lemont], 6:311

Measurement
A comparison of footprint indexes calculated from ink and electronic footprints [Urry and Wearing], 4:203
A comparison of radiographic foot measurements taken in two different positions [Bryant], 5:234
The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74
Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251
Radiographic analysis of metatarsus primus elevatus: a preliminary study [Christman et al], 6:294
The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121
Reliability of navicular displacement measurement as a clinical indicator of foot posture [Vinicombe et al], 5:262

Menopause
Juxtamalleolar lipoma: painful menopausal lipoma [Lemont], 6:311

Metatarsal
Brachymetatarsia of the third and fourth metatarsals [Goforth and Overbeek], 7:373
Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251
Radiographic analysis of metatarsus primus elevatus: a preliminary study [Christman et al], 6:294

Metatarsus primus elevatus
Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251
Radiographic analysis of metatarsus primus elevatus: a preliminary study [Christman et al], 6:294

Minority populations
Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313

Mycosis
The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456
Efficacy of terbinafine for toenail onychomycosis: a multicenter trial of various treatment durations [Pollak and Billstein], 3:127; (letter) [Wallace], 10:542
Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351
Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325
Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540

N

Nail
The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456
Efficacy of terbinafine for toenail onychomycosis: a multicenter trial of various treatment durations [Pollak and Billstein], 3:127; (letter) [Wallace], 10:542
Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351
Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325
Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540
Wound healing and infection in nail matrix phenolization wounds: does topical medication make a difference? [Dovison and Keenan], 5:230

Navicular
Reliability of navicular displacement measurement as a clinical indicator of foot posture [Vinicombe et al], 5:262

Necrobacillosis
Bilateral forefoot gangrene secondary to Lemierre's disease [Soave and Kuchar], 3:147

Neoplasm
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Hemangioma of the foot [Lamm and Pontious], 2:96
Juxtamalleolar lipoma: painful menopausal lipoma [Lemont], 6:311
Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109
Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79
Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93

Nerve
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451
Evaluation of pressure threshold prior to foot ulceration: one-*versus* two-point static touch [Barber et al], 10:508
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109
Surgically induced Charcot's foot [Fishco], 8:388
Syndrome of painful legs and moving toes [Yoon et al], 7:361

Neuroarthropathy
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365

Neuroma
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109

Neuropathy
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451
Evaluation of pressure threshold prior to foot ulceration: one-*versus* two-point static touch [Barber et al], 10:508
Surgically induced Charcot's foot [Fishco], 8:388

O

Onychomycosis
The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456
Efficacy of terbinafine for toenail onychomycosis: a multicenter trial of various treatment durations [Pollak and Billstein], 3:127; (letter) [Wallace], 10:542
Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351
Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325
Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540

Orthoses
Changes in foot function with the use of Root and Blake orthoses [Reed and Bennett], 4:184
The effect of customized insoles on the reduction of post-work discomfort [Sobel et al], 10:515
Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Foot orthosis prescription habits of Australian and New Zealand podiatric physicians [Landorf et al], 4:174
Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34

Osteomyelitis
Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365
The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445
Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369

Osteoporosis
Stress fracture of the foot secondary to osteoporosis: an atypical presentation [Freeman and Randall], 2:99

P

Pain
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Differentiation and treatment of phantom sensation, phantom pain, and residual-limb pain [Finnoff], 1:23
The effect of customized insoles on the reduction of post-work discomfort [Sobel et al], 10:515
The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Foot pain impairs balance and functional ability in community-dwelling older people [Menz and Lord], 5:222
Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356
Syndrome of painful legs and moving toes: a case study [Yoon et al], 7:361

Painful legs and moving toes syndrome
Syndrome of painful legs and moving toes: a case study [Yoon et al], 7:361

Pediatric population
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394

Periodontal disease
Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313

Peripheral vascular disease
Primary-care interventions for the population at risk for amputation [Yacyshyn and Andrews], 1:2

Pes planovalgus
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394

Phalanges
Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331

Phantom sensation
Differentiation and treatment of phantom sensation, phantom pain, and residual-limb pain [Finnoff], 1:23

Phenolization
Wound healing and infection in nail matrix phenolization wounds: does topical medication make a difference? [Dovison and Keenan], 5:230

Plantar fasciitis
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356
Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142

Poison
Voodoo powder as an etiology of leg ulcers [Reed and Wilson], 6:324

President's Message
APMA and the World Wide Web, 2:107
APMA's legislative agenda and advocacy, 7:385
Competency-based residency training, 5:273
Corporate partners, 6:329
e-Advocacy: your immediate link to members of Congress, 9:499
Fighting to level the playing field with managed care, 8:443
Medicare compliance for the podiatric medical practice, 4:219
Perspectives, 10:558
Public relations, 3:161
Why APMA?, 1:53

Pressure, plantar
Changes in foot function with the use of Root and Blake orthoses [Reed and Bennett], 4:184
The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280
Evaluation of pressure threshold prior to foot ulceration: one-*versus* two-point static touch [Barber et al], 10:508
Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275
Reduction of peak pressure on the forefoot with a rigid rocker-bottom postoperative shoe [Fuller et al], 10:501
Symmetry of plantar pressures and vertical forces in healthy subjects during walking [VanZant et al], 7:337

Pronation
The effect of high-Dye and low-Dye taping on rearfoot motion [Keenan and Tanner], 5:255
Function of the windlass mechanism in excessively pronated feet [Aquino and Payne], 5:245

Prostheses
Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34

Publication
Publication patterns and perceptions of the Australian podiatric medical faculty [Menz], 4:210

Public health
A conceptual model for public health education in podiatric medicine [Helfand and Hausman], 9:488
Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313

Q

Quality of life
Health-related quality of life following podiatric surgery [Bennett et al], 4:164

R

Radiographs
A comparison of radiographic foot measurements taken in two different positions [Bryant], 5:234
Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251
Radiographic analysis of metatarsus primus elevatus: a preliminary study [Christman et al], 6:294

Rearfoot
The effect of high-Dye and low-Dye taping on rearfoot motion [Keenan and Tanner], 5:255
The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121

Rehabilitation
Rehabilitation after amputation [Esquenazi and DiGiacomo], 1:13

Relaxation training
Effect of biofeedback-assisted relaxation training on foot ulcer healing [Rice et al], 3:132

Religion
Voodoo powder as an etiology of leg ulcers [Reed and Wilson], 6:324

Research
The case for multiple database searching in podiatric medicine (commentary) [Menz], 2:103
The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials [Moher et al], 8:437
Ensuring adequate reporting of randomized, controlled trials: the CONSORT statement (commentary) [Landorf], 8:435
Health-related quality of life following podiatric surgery [Bennett et al], 4:164
Publication patterns and perceptions of the Australian podiatric medical faculty [Menz], 4:210

Revascularization
The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533

S

Septicemia, postanginal
Bilateral forefoot gangrene secondary to Lemierre's disease [Soave and Kuchar], 3:147

Sesamoid
Internal fixation of a displaced tibial sesamoid fracture [Riley and Selner], 10:536
Isolated gouty tophus of a tibial sesamoid of the foot [Lemont and Sabo], 7:379

Skin
Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288
Evaluation of pressure threshold prior to foot ulceration: one-*versus* two-point static touch [Barber et al], 10:508
Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114
Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275
Voodoo powder as an etiology of leg ulcers [Reed and Wilson], 6:324

Smith-Lemli-Opitz syndrome
Smith-Lemli-Opitz syndrome: a genetic disorder with pedal manifestations [Sheff], 3:149

Splinting
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356

Sports medicine
Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
Podiatric medical applications of posterior night stretch splinting [Evans], 7:356
Survey of the effects of aerobic dance on the lower extremity in aerobic instructors [du Toit and Smith], 10:528

Spur
Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142

STA-peg
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394

Stability, postural
Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
Foot pain impairs balance and functional ability in community-dwelling older people [Menz and Lord], 5:222

Surgery
Arthrodesis of the interphalangeal joint of the hallux: a simple and effective technique [Yu et al], 8:427
Brachymetatarsia of the third and fourth metatarsals [Goforth and Overbeek], 7:373
Differentiation and treatment of phantom sensation, phantom pain, and residual-limb pain [Finnoff], 1:23
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394
Health-related quality of life following podiatric surgery [Bennett et al], 4:164
Internal fixation of a displaced tibial sesamoid fracture [Riley and Selner], 10:536
Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109
Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369
Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63
Primary-care interventions for the population at risk for amputation [Yacyshyn and Andrews], 1:2
Reduction of peak pressure on the forefoot with a rigid rocker-bottom postoperative shoe [Fuller et al], 10:501
Rehabilitation after amputation [Esquenazi and DiGiacomo], 1:13
Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142
The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533
Subtotal pedal amputations: biomechanical and intraoperative considerations [Funk and Young], 1:6
Surgically induced Charcot's foot [Fishco], 8:388
Talar fractures: three case studies [Jimenez and Morgan], 8:415
Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34
Wound healing and infection in nail matrix phenolization wounds: does topical medication make a difference? [Dovison and Keenan], 5:230

T

Talus
Posterior facet talocalcaneal coalition [Brekke et al], 8:422

Talar fractures: three case studies [Jimenez and Morgan], 8:415

Taping
The effect of high-Dye and low-Dye taping on rearfoot motion [Keenan and Tanner], 5:255

Tarsal tunnel syndrome
Evaluation of pressure threshold prior to foot ulceration: one-*versus* two-point static touch [Barber et al], 10:508
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109

Tarsus
Evaluation of pressure threshold prior to foot ulceration: one-*versus* two-point static touch [Barber et al], 10:508
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109
Posterior facet talocalcaneal coalition [Brekke et al], 8:422

Taste
Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540

Terbinafine
Efficacy of terbinafine for toenail onychomycosis: a multicenter trial of various treatment durations [Pollak and Billstein], 3:127; (letter) [Wallace], 10:542
Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540

Theory
Subtalar joint axis location and rotational equilibrium theory of foot function [Kirby], 9:465

Tibia
Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306

Tophus
Isolated gouty tophus of a tibial sesamoid of the foot [Lemont and Sabo], 7:379

Tumor
Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Hemangioma of the foot [Lamm and Pontious], 2:96
Juxtamalleolar lipoma: painful menopausal lipoma [Lemont], 6:311
Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306
Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109
Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79
Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93

U

Ulcer
Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451
The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280
Effect of biofeedback-assisted relaxation training on foot ulcer healing [Rice et al], 3:132
Evaluation of pressure threshold prior to foot ulceration: one-*versus* two-point static touch [Barber et al], 10:508
Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275
Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369
A prospective study of predictors for foot ulceration in type 2 diabetes [Kästenbauer et al], 7:343
Voodoo powder as an etiology of leg ulcers [Reed and Wilson], 6:324
Wound healing and infection in nail matrix phenolization wounds: does topical medication make a difference? [Dovison and Keenan], 5:230

V

Vascular disease
Primary-care interventions for the population at risk for amputation [Yacyshyn and Andrews], 1:2

Vein
Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93

Verrucae
Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79

Vohwinkel's syndrome
Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114

Voodoo
Voodoo powder as an etiology of leg ulcers [Reed and Wilson], 6:324

W

Warts
Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79

Windlass mechanism
Function of the windlass mechanism in excessively pronated feet [Aquino and Payne], 5:245

Work
The effect of customized insoles on the reduction of post-work discomfort [Sobel et al], 10:515

AUTHOR INDEX

A

Abu-Rumman, Patricia L.: Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451

Altman, Douglas G.: The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials [Moher et al], 8:437

Anarella, James J.: Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325

Andrews, Karen L.: Primary-care interventions for the population at risk for amputation [Yacyshyn and Andrews], 1:2

Aquino, Anna: Function of the windlass mechanism in excessively pronated feet [Aquino and Payne], 5:245

Armstrong, David G.:

Activity monitors: should we begin dosing activity as we dose a drug? [Armstrong and Boulton], 3:152

Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451

Auinger, M.: A prospective study of predictors for foot ulceration in type 2 diabetes [Kästenbauer et al], 7:343

B

Bakotic, Bradley: Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351

Barber, Mitchell A.: Evaluation of pressure threshold prior to foot ulceration: one- *versus* two-point static touch [Barber et al], 10:508

Barbosa, Peter: Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79

Bauer, Gary R.:

Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331

Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63

Beegle, Terrance: Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89

Bennett, Paul J.:

Changes in foot function with the use of Root and Blake orthoses [Reed and Bennett], 4:184

Health-related quality of life following podiatric surgery [Bennett et al], 4:164

Bernard, Joshua M.: Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306

Billstein, Stephan A.: Efficacy of terbinafine for toenail onychomycosis: a multicenter trial of various treatment durations [Pollak and Billstein], 3:127

Black, James A.: Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306

Boc, Steven F.: Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365

Borkowski, Piotr: Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351

Botek, Georgeanne: Syndrome of painful legs and moving toes: a case study [Yoon et al], 7:361

Boulton, Andrew J. M.:

Activity monitors: should we begin dosing activity as we dose a drug? [Armstrong and Boulton], 3:152

Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451

Bowker, Peter: Scientific approach to the axis of rotation at the midtarsal joint [Nester et al], 2:68

Brazzo, Keith: Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365

Brekke, Mark K.: Posterior facet talocalcaneal coalition [Brekke et al], 8:422

Broering, Jeanette M.: Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521

Brook, Joel W.: Arthrodesis of the interphalangeal joint of the hallux: a simple and effective technique [Yu et al], 8:427

Bryant, Alan: Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251

Bryant, Jennifer A.: A comparison of radiographic foot measurements taken in two different positions, 5:234

C

Caldwell, Bryan D.: Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114

Campion, Nicholas: The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

Caselli, Mark A.: The effect of customized insoles on the reduction of postwork discomfort [Sobel et al], 10:515

Chang, Thomas J.: Hemophilic arthropathy: considerations in management [Chang et al], 8:406

Christman, Robert A.: Radiographic analysis of metatarsus primus elevatus: a preliminary study [Christman et al], 6:294

Christos, Paul J.: The effect of customized insoles on the reduction of postwork discomfort [Sobel et al], 10:515

Conolley, Janice: Evaluation of pressure threshold prior to foot ulceration: one- *versus* two-point static touch [Barber et al], 10:508

Cornwall, Mark W.:

The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280

Symmetry of plantar pressures and vertical forces in healthy subjects during walking [VanZant et al], 7:337

Crabtree, Christopher: Syndrome of painful legs and moving toes: a case study [Yoon et al], 7:361

D

Dananberg, Howard J.: Manipulation method for the treatment of ankle equinus (letter), 2:105

Datta, Ujjwal K.: The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

DeBello, John A.: Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325

Dellon, A. Lee:

Evaluation of pressure threshold prior to foot ulceration: one- *versus* two-point static touch [Barber et al], 10:508

Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109

DiGiacomo, Robert: Rehabilitation after amputation [Esquenazi and DiGiacomo], 1:13
DiLandro, Anthony C.: The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300
Dixon, Russell M.: Effect of biofeedback-assisted relaxation training on foot ulcer healing [Rice et al], 3:132
Dovison, Russell: Wound healing and infection in nail matrix phenolization wounds: does topical medication make a difference? [Dovison and Keenan], 5:230
Doyle, Joseph J.: Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Dunne, Michael P.: Health-related quality of life following podiatric surgery [Bennett et al], 4:164
du Toit, Verona: Survey of the effects of aerobic dance on the lower extremity in aerobic instructors [du Toit and Smith], 10:528

E

Edwards, Jenifer: Reduction of peak pressure on the forefoot with a rigid rocker-bottom postoperative shoe [Fuller et al], 10:501
Esquenazi, Alberto: Rehabilitation after amputation [Esquenazi and DiGiacomo], 1:13
Evans, Angela: Podiatric medical applications of posterior night stretch splinting, 7:356

F

Feldman, Kent: Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394
Fikar, Charles R.: Internet resources for podiatric medical students [Fikar and Fikar], 6:316
Fikar, Ronald M.: Internet resources for podiatric medical students [Fikar and Fikar], 6:316
Findlow, Andrew: Scientific approach to the axis of rotation at the midtarsal joint [Nester et al], 2:68
Finnoff, Jonathan: Differentiation and treatment of phantom sensation, phantom pain, and residual-limb pain, 1:23
Fiorilli, Anthony: Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331
Fishco, William D.: Surgically induced Charcot's foot, 8:388
Fisher, Susan Gross: Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275
Flake, Edward: Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394
Flanigan, K. Paul: Radiographic analysis of metatarsus primus elevatus: a preliminary study [Christman et al], 6:294
Fleischli, Jeffrey W.: The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445
Forg, Patricia: Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394
Freeman, Denise: Stress fracture of the foot secondary to osteoporosis: an atypical presentation [Freeman and Randall], 2:99
Fuller, Eric: Reduction of peak pressure on the forefoot with a rigid rocker-bottom postoperative shoe [Fuller et al], 10:501
Funk, Christopher: Subtotal pedal amputations: biomechanical and intraoperative considerations [Funk and Young], 1:6

G

Gaggero, Geoffrey L.: Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79
Garrow, Adam P.: The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74
Gause, Douglas: Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Giorgini, Renato J.: Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34
Goforth, William P.:
Brachymetatarsia of the third and fourth metatarsals [Goforth and Overbeek], 7:373
Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55
Grandi, Robert: Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93
Green, Donald R.:
Flake-Austin modification of the STA-peg arthroereisis: a retrospective study [Forg et al], 8:394
Posterior facet talocalcaneal coalition [Brekke et al], 8:422

H

Hambleton, Julie: Hemophilic arthropathy: considerations in management [Chang et al], 8:406
Handzel, Charleen: Isolated cuboid fracture: a rare occurrence [Miller and Handzel], 2:85
Harkless, Lawrence B.: The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533
Hausman, Alice J.: A conceptual model for public health education in podiatric medicine [Helfand and Hausman], 9:488
Hayes, Walter W.: The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456
Helfand, Arthur E.: A conceptual model for public health education in podiatric medicine [Helfand and Hausman], 9:488
Hillstrom, Howard J.:
Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331
Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63
The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121
Hodor, Lawrence: Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369
Holewinski, Jean: Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351
Holland, Paul: Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313
Hosch, Jonathan C.: Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55
Hughes, Monica L.: Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114
Hughes, Thomas J.: Molecular quantification of human β-glucuronidase levels in a patient with Vohwinkel's syndrome [Hughes et al], 3:114
Husain, Khalid: The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533

I

Irsigler, K.: A prospective study of predictors for foot ulceration in type 2 diabetes [Kästenbauer et al], 7:343

J

Japour, Christopher J.: Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34

Jayson, Malcolm I. V.: The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74

Jimenez, A. Louis: Talar fractures: three case studies [Jimenez and Morgan], 8:415

Joseph, Warren S.:

Guidelines for authors, 1:50, 5:270, 7:382, 9:496

Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521

K

Kaleta, Jennifer L.: The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445

Kalker, Alan J.: Effect of biofeedback-assisted relaxation training on foot ulcer healing [Rice et al], 3:132

Karr, Jeffrey C.: Magnetic resonance imaging evaluation of monostotic fibrous dysplasia of the tibia [Karr et al], 6:306

Kästenbauer, T.: A prospective study of predictors for foot ulceration in type 2 diabetes [Kästenbauer et al], 7:343

Kaufman, Jeffrey L.: Bilateral forefoot gangrene secondary to Lemierre's disease (letter), 7:381

Keenan, Anne-Maree:

The effect of high-Dye and low-Dye taping on rearfoot motion [Keenan and Tanner], 5:255

Foot orthosis prescription habits of Australian and New Zealand podiatric physicians [Landorf et al], 4:174

Wound healing and infection in nail matrix phenolization wounds: does topical medication make a difference? [Dovison and Keenan], 5:230

Kim, Jaesuk: Pain at the site of tarsal tunnel incision due to neuroma of the posterior branch of the saphenous nerve [Kim and Dellon], 3:109

Kirby, Kevin A.: Subtalar joint axis location and rotational equilibrium theory of foot function, 9:465

Kuchar, David J.: Bilateral forefoot gangrene secondary to Lemierre's disease [Soave and Kuchar], 3:147

L

La Fontaine, Javier: The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533

Lamm, Bradley M.:

Hemangioma of the foot [Lamm and Pontious], 2:96

Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331

Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63

Landino, Thomas: Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365

Landorf, Karl B.:

Ensuring adequate reporting of randomized, controlled trials: the CONSORT statement (commentary), 8:435

Foot orthosis prescription habits of Australian and New Zealand podiatric physicians [Landorf et al], 4:174

Lane, George: Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93

LaPointe, Stephan J.: The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121

Lavian, Dana: Acute Charcot foot changes *versus* osteomyelitis: does Tc-99m HMPAO Labeled Leukocytes Scan differentiate? [Boc et al], 7:365

Lehmann, Sean: Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369

Lemont, Harvey:

Isolated gouty tophus of a tibial sesamoid of the foot [Lemont and Sabo], 7:379

Juxtamalleolar lipoma: painful menopausal lipoma, 6:311

Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540

Lepore, Frank Louis: The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

Levitz, Steven J.:

The effect of customized insoles on the reduction of post-work discomfort [Sobel et al], 10:515

Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34

Lieberman, Ron: Posterior facet talocalcaneal coalition [Brekke et al], 8:422

Lilja, Erik C.: The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300

Lord, Stephen R.: Foot pain impairs balance and functional ability in community-dwelling older people [Menz and Lord], 5:222

Loveland, Lacey: Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79

Lynch, D. Matt: Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55

M

Macfarlane, Gary J.: The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74

Mahoney, Belinda: Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251

Martin, Joe E.: Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55

McCreary, Jon P.: The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456

McPoil, Thomas G.:

The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280

Symmetry of plantar pressures and vertical forces in healthy subjects during walking [VanZant et al], 7:337

Menz, Hylton B.:

The case for multiple database searching in podiatric medicine (commentary), 2:103

Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194

Foot pain impairs balance and functional ability in community-dwelling older people [Menz and Lord], 5:222

Manipulation method for the treatment of ankle equinus (letter), 2:105

Publication patterns and perceptions of the Australian podiatric medical faculty, 4:210

Reliability of navicular displacement measurement as a clinical indicator of foot posture [Vinicombe et al], 5:262

Miller, S. Rick: Isolated cuboid fracture: a rare occurrence [Miller and Handzel], 2:85

Miller, Stephen J.:

Review of *Functional Reconstruction of the Foot and Ankle*, 5:269

Review of *Handbook of Lower Extremity Neurology*, 2:102

Review of *Textbook of Bunion Surgery*, 5:269

Moadab, Anoosh: Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288

Mohamed, Shenin: Hemophilic arthropathy: considerations in management [Chang et al], 8:406
Moher, David: The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials [Moher et al], 8:437
Morgan, James H., Jr.: Talar fractures: three case studies [Jimenez and Morgan], 8:415
Murff, Randall T.: Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55
Murphy, R. Dan: Partial calcanectomy in the treatment of chronic heel ulceration [Lehmann et al], 7:369

N

Nakra, Aprajita: The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121
Nester, Christopher J.: Scientific approach to the axis of rotation at the midtarsal joint [Nester et al], 2:68
Ng, Alan: Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Nixon, Brent P.: Continuous activity monitoring in persons at high risk for diabetes-related lower-extremity amputation [Armstrong et al], 9:451
Noriega, James A.: Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142

O

Odom, Richard D.: Mechanical treatment of plantar fasciitis: a prospective study [Martin et al], 2:55
Overbeek, Troy D.: Brachymetatarsia of the third and fourth metatarsals [Goforth and Overbeek], 7:373

P

Papageorgiou, Ann: The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74
Patterson, Carla: Health-related quality of life following podiatric surgery [Bennett et al], 4:164
Payne, Craig: Function of the windlass mechanism in excessively pronated feet [Aquino and Payne], 5:245
Peebles, Charles: The reliability of clinical and caliper-based calcaneal bisection measurements [LaPointe et al], 3:121
Percy, Matthew L.: Effects of prefabricated foot orthoses and soft insoles on postural stability in professional soccer players [Percy and Menz], 4:194
Pollak, Richard A.: Efficacy of terbinafine for toenail onychomycosis: a multicenter trial of various treatment durations [Pollak and Billstein], 3:127; (letter), 10:542
Pontious, Jane: Hemangioma of the foot [Lamm and Pontious], 2:96
Pozarny, Edward: Venous aneurysm of the dorsal venous arch [Pozarny et al], 2:93

R

Randall, David B.: Stress fracture of the foot secondary to osteoporosis: an atypical presentation [Freeman and Randall], 2:99
Raspovic, Anita: Reliability of navicular displacement measurement as a clinical indicator of foot posture [Vinicombe et al], 5:262
Reed, Bradley: Voodoo powder as an etiology of leg ulcers [Reed and Wilson], 6:324
Reed, Lloyd: Changes in foot function with the use of Root and Blake orthoses [Reed and Bennett], 4:184
Reilly, Charles H.: The diagnosis of osteomyelitis in diabetes using erythrocyte sedimentation rate: a pilot study [Kaleta et al], 9:445
Reyzelman, Alex: The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533
Ribeiro, Carla E.:
Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331
Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63
Rice, Birgitta: Effect of biofeedback-assisted relaxation training on foot ulcer healing [Rice et al], 3:132
Richardson, Hugh L.: Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34
Riley, Jay: Internal fixation of a displaced tibial sesamoid fracture [Riley and Selner], 10:536
Robbins, Jeffrey M.: Passage of policy statement on the high risk of blindness, lower-extremity amputations, and oral health consequences in minority populations due to diabetes [Robbins and Holland], 6:313
Rockett, Andrea K.: Atypical presentation of plantar fasciitis secondary to soft-tissue mass infiltration [Ng et al], 2:89
Rosenblum, Jonathan: The effect of customized insoles on the reduction of postwork discomfort [Sobel et al], 10:515
Rothenberg, Gary: The role of revascularization in transmetatarsal amputations [La Fontaine et al], 10:533
Rupley, Kathryne F.: Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288
Rushworth, R. Louise: Foot orthosis prescription habits of Australian and New Zealand podiatric physicians [Landorf et al], 4:174

S

Sabo, Marc:
Isolated gouty tophus of a tibial sesamoid of the foot [Lemont and Sabo], 7:379
Terbinafine-associated taste disturbance with normal taste threshold scores [Lemont and Sabo], 10:540
Sage, Ronald A.: Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275
Sauseng, S.: A prospective study of predictors for foot ulceration in type 2 diabetes [Kästenbauer et al], 7:343
Schein, Jeffrey R.: Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Scherer, William P.: The diagnosis of onychomycosis in a geriatric population: a study of 450 cases in South Florida [Scherer et al], 9:456
Schindler, Jay V.: Effect of biofeedback-assisted relaxation training on foot ulcer healing [Rice et al], 3:132
Schroeder, Stephen: Reduction of peak pressure on the forefoot with a rigid rocker-bottom postoperative shoe [Fuller et al], 10:501
Schulz, Kenneth F.: The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials [Moher et al], 8:437
Segura, Lolita: Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79
Selner, Marc: Internal fixation of a displaced tibial sesamoid fracture [Riley and Selner], 10:536
Sheff, Jordan S.: Smith-Lemli-Opitz syndrome: a genetic disorder with pedal manifestations, 3:149
Signorile, Joseph: The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300
Silman, Alan J.: The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74

Slattery, Matthew: The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Smith, Richard: Survey of the effects of aerobic dance on the lower extremity in aerobic instructors [du Toit and Smith], 10:528
Smith, Wayne: The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280
Smith, William K.: Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142
Smith, William K., Jr.: Resection of a plantar calcaneal spur using the holmium:yttrium-aluminum-garnet (Ho:YAG) laser [Smith et al], 3:142
Soave, Ronald L.: Bilateral forefoot gangrene secondary to Lemierre's disease [Soave and Kuchar], 3:147
Sobel, Ellen:
The effect of customized insoles on the reduction of post-work discomfort [Sobel et al], 10:515
Use of prostheses and footwear in 110 inner-city partial-foot amputees [Sobel et al], 1:34
Sokol, G.: A prospective study of predictors for foot ulceration in type 2 diabetes [Kästenbauer et al], 7:343
Sorrento, Dean L.: Radiographic analysis of metatarsus primus elevatus: a preliminary study [Christman et al], 6:294
Sowell, Robert D. "Doug":
APMA and the World Wide Web, 2:107
APMA's legislative agenda and advocacy, 7:385
Competency-based residency training, 5:273
Corporate partners, 6:329
e-Advocacy: your immediate link to members of Congress, 9:499
Fighting to level the playing field with managed care, 8:443
Medicare compliance for the podiatric medical practice, 4:219
Perspectives, 10:558
Public relations, 3:161
Why APMA?, 1:53
Spaulding, Cecily M.: Evaluation of pressure threshold prior to foot ulceration: one- *versus* two-point static touch [Barber et al], 10:508
Stanich, Christopher C.: Radiographic analysis of metatarsus primus elevatus: a preliminary study [Christman et al], 6:294
Stier, David M.: Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521

T

Tanner, Craig M.: The effect of high-Dye and low-Dye taping on rearfoot motion [Keenan and Tanner], 5:255
Thomas, Elaine: The grading of hallux valgus: the Manchester scale [Garrow et al], 2:74
Tinley, Paul:
The efficacy of functional foot orthoses in the control of pain in ankle joint disintegration in hemophilia [Slattery and Tinley], 5:240
Lateral intermetatarsal angle: a useful measurement of metatarsus primus elevatus? [Bryant et al], 5:251
Toth, Christopher: Preventing complications in the diabetic patient with toenail onychomycosis [Anarella et al], 6:325

U

Urry, Stephen R.: A comparison of footprint indexes calculated from ink and electronic footprints [Urry and Wearing], 4:203

V

VanZant, R. Scott: Symmetry of plantar pressures and vertical forces in healthy subjects during walking [VanZant et al], 7:337
Vargo, Frank E.: Arthrodesis of the interphalangeal joint of the hallux: a simple and effective technique [Yu et al], 8:427
Vinicombe, Ann: Reliability of navicular displacement measurement as a clinical indicator of foot posture [Vinicombe et al], 5:262
Viscovich, John B.: The prevalence of the arcuate artery: a cadaveric study of 72 feet [DiLandro et al], 6:300
Vlahovic, Tracey C.:
Lesser proximal interphalangeal joint arthrodesis: a retrospective analysis of the peg-in-hole and end-to-end procedures [Lamm et al], 7:331
Peg-in-hole, end-to-end, and V arthrodesis: a comparison of digital stabilization in fresh cadaveric specimens [Lamm et al], 2:63

W

Wadhams, Peter: Effectiveness of a nonrinse, alcohol-free antiseptic hand wash [Moadab et al], 6:288
Wallace, George F.: Efficacy of terbinafine for toenail onychomycosis (letter), 10:542
Warolin, Karen L.: Patient satisfaction with oral *versus* nonoral therapeutic approaches in onychomycosis [Stier et al], 10:521
Wearing, Scott C.: A comparison of footprint indexes calculated from ink and electronic footprints [Urry and Wearing], 4:203
Webster, Julie Kate: Outpatient care and morbidity reduction in diabetic foot ulcers associated with chronic pressure callus [Sage et al], 6:275
Whitaker, Jeffrey M.: Plantar verrucae in patients with human immunodeficiency virus: clinical presentation and treatment response [Whitaker et al], 2:79
Williams, Marie: Onychomycosis: an analysis of 50 cases and a comparison of diagnostic techniques [Borkowski et al], 7:351
Wilson, Richard C.: Voodoo powder as an etiology of leg ulcers [Reed and Wilson], 6:324
Wright, Ethel: Posterior facet talocalcaneal coalition [Brekke et al], 8:422

Y

Yacyshyn, Vincent J.: Primary-care interventions for the population at risk for amputation [Yacyshyn and Andrews], 1:2
Yamada, Wesley: The distribution of plantar pressures in American Indians with diabetes mellitus [McPoil et al], 6:280
Yoon, Joonhyun: Syndrome of painful legs and moving toes: a case study [Yoon et al], 7:361
Young, Gregg: Subtotal pedal amputations: biomechanical and intraoperative considerations [Funk and Young], 1:6
Yu, Gerard V.: Arthrodesis of the interphalangeal joint of the hallux: a simple and effective technique [Yu et al], 8:427